

Perturbative pion form factor reexamined

David Tung and Hsiang-nan Li

Department of Physics, National Chung-Cheng University,
Chia-Yi, Taiwan, Republic of China

PACS: 13.40.Gp, 12.38.Bx, 12.38.Cy

Abstract

We recalculate the pion electromagnetic form factor based on the perturbative QCD formalism that includes the Sudakov resummation. We take into account the evolution of the pion wave function in b , which represents the transverse extent of the pion. An infrared enhancement is observed when $1/b$ approaches Λ_{QCD} . We propose to freeze the evolution of the wave function at some scale above Λ_{QCD} , such that perturbative contributions are stabilized. Our predictions are consistent with experimental data, and insensitive to the variation of relevant parameters.

I. Introduction

The applicability of perturbative QCD (PQCD) to exclusive processes has been a controversy for almost two decades [1, 2]. Although there is a general agreement that PQCD can successfully make predictions for exclusive reactions as momentum transfers go to infinity, it remains unclear whether experimentally accessible energy scales are large enough to justify these predictions. A progress was made recently by Li and Sterman [3], who proposed a modified PQCD formalism with Sudakov suppression taken into account. They found that nonperturbative contributions are suppressed significantly by the Sudakov effects, and the applicability of PQCD can be extended down to few GeV scales. However, their analysis of the pion electromagnetic form factor did not concern the match with experimental data, such that a convincing piece of justification of their approach is missing. In this paper we shall further include the evolution of the pion wave function into the modified PQCD formalism, and demonstrate that our predictions explain the data.

The PQCD theory for exclusive processes was first proposed by Brodsky and Lepage [4]. They argued that an exclusive process, such as the pion form factor, can be factorized into two types of subprocesses: wave functions which carry the nonperturbative information of the initial and final state pions, and a hard amplitude which describes the scattering of a valence quark of the pion off the energetic photon. The former is not calculable in perturbation theory, and needs to be parametrized by a model, to be derived by nonperturbative methods, such as QCD sum rules and lattice gauge theory, or to be determined by experimental data. The latter, characterized by a large momentum flow, is calculable in perturbation theory. Combining these two types of subprocesses, predictions for the pion form factor at experimentally accessible energies scales can be made.

According to this naive picture, the factorization formula for the pion form factor $F_\pi(Q^2)$, graphically represented by Fig. 1(a), is written as

$$F_\pi(Q^2) = \int_0^1 dx_1 dx_2 \phi(x_2, \mu) H(x_1, x_2, Q, \mu) \phi(x_1, \mu) . \quad (1)$$

$Q^2 = -2P_1 \cdot P_2$ is the momentum transfer from the photon, P_1 (P_2) being the momentum of the incoming (outgoing) pion. μ is a renormalization and factorization scale, below which QCD dynamics is regarded as being non-perturbative and absorbed into the pion wave function ϕ , and above which

QCD dynamics is regarded as being perturbative and absorbed into the hard amplitude H . $\phi(x, \mu)$ gives the probability of a valence quark carrying a fractional momentum xP in the parton model at the scale μ . Usually, the μ dependence of ϕ is neglected, *i.e.*, $\phi(x, \mu) = \phi(x)$. H is obtained by computing quark-photon scattering diagrams.

To make predictions for F_π , one substitutes the lowest-order expression of H derived from Fig. 1(b),

$$H(x_1, x_2, Q, \mu) = \frac{16\pi C_F \alpha_s(\mu)}{x_1 x_2 Q^2}, \quad (2)$$

and the asymptotic pion wave function [5]

$$\phi^{AS}(x) = \frac{3f_\pi}{\sqrt{2N_c}} x(1-x), \quad (3)$$

into Eq. (1), where $\alpha_s(\mu)$ is the running coupling constant, $C_F = 4/3$ the color factor, $N_c = 3$ the number of colors, and $f_\pi = 0.133$ GeV the pion decay constant. Because ϕ^{AS} peaks at $x = 1/2$, the main contributions to Eq. (1) come from the region with intermediate x . Higher-order corrections to H then produce logarithms of the type $\ln(Q^2/\mu^2)$, which may be so large as to spoil the perturbative expansion. To eliminate the logarithms, a natural choice of μ is $\mu = Q$. It is easy to find that $Q^2 F_\pi(Q^2) \sim 0.12$ GeV² for $Q^2 \sim 4$ -10 GeV² is only 1/3 of the data ~ 0.35 GeV² [6, 7, 8, 9]. This contradiction implies that the pion form factor has not yet become completely asymptotic at experimentally accessible energies.

Hence, one may resort to a preasymptotic model, the Chernyak and Zhitnitsky (CZ) wave function derived from QCD sum rules [10],

$$\phi^{CZ}(x) = \frac{15f_\pi}{\sqrt{2N_c}} x(1-x)(1-2x)^2, \quad (4)$$

which possesses maxima at the end points $x \rightarrow 0$ and $x \rightarrow 1$. The CZ wave function enhances the predictions for $Q^2 F_\pi(Q^2)$ to 0.30 GeV², and improves the match with the data. However, such a success was criticized by Isgur and Llewellyn Smith [1, 11] and by Radyushkin [12]: Since ϕ^{CZ} emphasizes the contributions from small x , higher-order corrections to H in fact produce logarithms like $\ln(x_1 x_2 Q^2/\mu^2)$. Hence, the natural choice of μ is

$\mu = \sqrt{x_1 x_2} Q$, which is consistent with $\mu = Q$ employed above for intermediate x , but results in a large coupling constant $\alpha_s(\sqrt{x_1 x_2} Q) > 1$ at small x , indicating that the end-point regions are nonperturbative regions. A careful analysis revealed that the enhancement of the predictions for $Q^2 F_\pi(Q^2)$ is due to the amplification of the nonperturbative end-point contributions. As a consequence, the perturbative calculation loses its self-consistency as a weak-coupling expansion.

The above discussions hint that the end-point regions should be treated in a different way. In general, a valence quark in the pion can carry a small amount of transverse momenta k_T , which then flow through the hard scattering. Therefore, the hard gluon propagator is written as

$$\frac{1}{x_1 x_2 Q^2 + k_T^2} \approx \frac{1}{x_1 x_2 Q^2} \left(1 - \frac{k_T^2}{x_1 x_2 Q^2} \right), \quad (5)$$

where the first term corresponds exactly to the asymptotic H in Eq. (2), and the second is suppressed by powers of Q^2 at intermediate x . If the end-point regions are not important, one may drop the second term, and integrate out the k_T dependence in the wave functions, arriving at the leading-power factorization formula in Eq. (1). However, the end-point difficulties indicate that the higher-power effects are crucial, and should be kept at the outset as deriving the perturbative expression for the pion form factor.

The introduction of k_T leads to three observations immediately. First, the pion form factor becomes a two-scale (Q and k_T) problem, and the resummation technique is required to organize the large logarithms $\ln(Q^2/k_T^2)$ from radiative corrections to the wave functions. Second, the process must be analyzed in the Fourier transform space of k_T , denoted by b [13], which is regarded as the transverse separation between the valence quarks of the pion. Third, the quantity $1/b$, now considered as one of the characteristic scales of the hard amplitude, should be substituted for the argument of α_s if $1/b > \sqrt{x_1 x_2} Q$.

Li and Sterman found that the resummation of the large logarithms suppresses the elastic scattering at large spatial separation. This property, called Sudakov suppression [13, 14, 15], makes nonperturbative contributions from large b , no matter what x are, less important, and improves the applicability of PQCD at few GeV scales. However, their discussion was restricted to the PQCD applicability, such that the results of the pion form factor do not

match the data. In the present work we shall aim to explain the experimental data by including a new piece of information, the evolution of the wave function, into the modified PQCD formalism. Our predictions are found to be dominated by perturbative contributions for $Q^2 > 4 \text{ GeV}^2$, and in good agreement with the data.

A brief review of the standard asymptotic expression for the form factors [5, 10, 16, 17, 18, 19] may help to motivate our viewpoint. The factorization formula for the pion form factor including the evolution of the wave function is derived in Sect. II. Numerical results are given in Sect. III. Section IV is the conclusion.

II. The Factorization Formula

In this section we review factorization theorems including Sudakov effects for the pion form factor, which is expressed as the convolution of a hard amplitude with wave functions. We investigate radiative corrections to the quark-photon scattering diagrams shown in Fig. 2, and explain how they are factorized into the convolution factors. There are two types of important corrections: collinear, when the loop momentum is parallel to the incoming or outgoing pion momentum, and soft, when the loop momentum is much smaller than the momentum transfer Q^2 . Each type of the important corrections gives rise to large single logarithms. They may overlap to produce double logarithms in some cases. In axial gauge the two-particle reducible diagrams, like Figs. 2(a) and 2(b), give double logarithms from the overlap of collinear and soft divergences, while the two-particle irreducible corrections, like Figs. 2(c) and 2(d), give only single soft logarithms.

It can be shown that soft divergences cancel between Figs. 2(a) and 2(b), as well as between 2(c) and 2(d), in the asymptotic region with $b \rightarrow 0$. Therefore, reducible corrections are dominated by collinear divergences, and can be absorbed into the pion wave function $\mathcal{P}$, which involves similar dynamics. Irreducible corrections, due to the cancellation of soft divergences, are then absorbed into the hard scattering amplitude H . Based on the above reasoning, the factorization formula for the pion form factor is written in the b

space as [3],

$$F_\pi(Q^2) = \int_0^1 dx_1 dx_2 \int \frac{d^2 \mathbf{b}}{(2\pi)^2} \mathcal{P}(x_2, b, P_2, \mu) \times \tilde{H}(x_1, x_2, b, Q, \mu) \mathcal{P}(x_1, b, P_1, \mu) , \quad (6)$$

where $\tilde{H}$ is the Fourier transform of H .

We choose the Breit frame such that $P_1^+ = P_2^- = Q/\sqrt{2}$ and all other components of P 's vanish. After resumming the double logarithms, we obtain

$$\mathcal{P}(x, b, P, \mu) = \exp[-s(x, b, Q) - s(1-x, b, Q)] \bar{\mathcal{P}}(x, b, \mu) . \quad (7)$$

The exponent $s(\xi, b, Q)$, $\xi = x$ and $1-x$, is expressed as [14]

$$s(\xi, b, Q) = \int_{1/b}^{\xi Q/\sqrt{2}} \frac{dp}{p} \left[\ln \left(\frac{\xi Q}{\sqrt{2}p} \right) A(\alpha_s(p)) + B(\alpha_s(p)) \right] , \quad (8)$$

where the anomalous dimensions A to two loops and B to one loop are given by

$$\begin{aligned} A &= \mathcal{C}_F \frac{\alpha_s}{\pi} + \left[\frac{67}{9} - \frac{\pi^2}{3} - \frac{10}{27} n_f + \frac{8}{3} \beta_1 \ln \left(\frac{e^{\gamma_E}}{2} \right) \right] \left(\frac{\alpha_s}{\pi} \right)^2 , \\ B &= \frac{2}{3} \frac{\alpha_s}{\pi} \ln \left(\frac{e^{2\gamma_E-1}}{2} \right) , \end{aligned} \quad (9)$$

γ_E being the Euler constant. The two-loop running coupling constant,

$$\frac{\alpha_s(\mu)}{\pi} = \frac{4}{\beta_0 \ln(\mu^2/\Lambda^2)} - \frac{16\beta_1}{\beta_0^3} \frac{\ln \ln(\mu^2/\Lambda^2)}{\ln^2(\mu^2/\Lambda^2)} , \quad (10)$$

with the coefficients

$$\beta_0 = \frac{33-2n_f}{3} , \quad \beta_1 = \frac{153-19n_f}{6} , \quad (11)$$

and the QCD scale $\Lambda \equiv \Lambda_{\text{QCD}}$, will be substituted into Eq. (8). We require the relation of the involved scales $\xi Q/\sqrt{2} > 1/b > \Lambda$ as indicated by the bounds of the variable p in Eq. (8). QCD dynamics below $1/b$ is regarded as being nonperturbative, and absorbed into the initial condition $\bar{\mathcal{P}}(x, b, \mu)$.

The functions $\bar{\mathcal{P}}$ and $\tilde{H}$ still contain single logarithms from ultraviolet divergences, which can be summed using the RG methods [14]. The large- b behavior of $\mathcal{P}$ is then summarized as

$$\begin{aligned} \mathcal{P}(x, b, P, \mu) &= \exp \left[-s(x, b, Q) - s(1-x, b, Q) - 2 \int_{1/b}^{\mu} \frac{d\bar{\mu}}{\bar{\mu}} \gamma_q(\alpha_s(\bar{\mu})) \right] \\ &\times \bar{\mathcal{P}}(x, b, 1/b) , \end{aligned} \quad (12)$$

with $\gamma_q = -\alpha_s/\pi$ the quark anomalous dimension in axial gauge. To concentrate on the evolution of $\bar{\mathcal{P}}$ in $1/b$, we ignore the intrinsic b dependence denoted by the argument b [20], that is, we adopt $\bar{\mathcal{P}}(x, b, 1/b) = \phi(x, 1/b)$.

Since the pion form factor, as a physical observable, is μ independent, the RG analysis applied to $\tilde{H}$ gives

$$\begin{aligned} \tilde{H}(x_i, b, Q, \mu) &= \exp \left[-4 \int_{\mu}^t \frac{d\bar{\mu}}{\bar{\mu}} \gamma_q(\alpha_s(\bar{\mu})) \right] \\ &\times \tilde{H}(x_i, b, Q, t) , \end{aligned} \quad (13)$$

where t is the largest mass scale involved in the hard scattering,

$$t = \max(\sqrt{x_1 x_2} Q, 1/b) . \quad (14)$$

The scale $\sqrt{x_1 x_2} Q$ is associated with the longitudinal momentum of the hard gluon and $1/b$ with the transverse momentum. At lowest order, H is given by

$$H(x_i, \mathbf{k}_{T_i}, Q) = \frac{16\pi\alpha_s\mathcal{C}_F x_1 Q^2}{(x_1 Q^2 + \mathbf{k}_{1T}^2)(x_1 x_2 Q^2 + (\mathbf{k}_{1T} - \mathbf{k}_{2T})^2)} \quad (15)$$

$$\sim \frac{16\pi\alpha_s\mathcal{C}_F}{x_1 x_2 Q^2 + (\mathbf{k}_{1T} - \mathbf{k}_{2T})^2} , \quad (16)$$

where we have neglected the transverse momentum in the numerator. In the second form, we have further neglected the transverse momentum associated with the virtual fermion lines in H , which are linear rather than quadratic in x . Because Eq. (16) depends on the combination of transverse momenta, $\tilde{H}$ involves only a single b , which has been made explicit in Eq. (6).

Combining all the above ingredients, we arrive at the factorization formula for the pion form factor,

$$\begin{aligned} F_{\pi}(Q^2) &= 16\pi\mathcal{C}_F \int_0^1 dx_1 dx_2 \int_0^{\infty} b db \phi(x_1, 1/b) \phi(x_2, 1/b) \alpha_s(t) K_0(\sqrt{x_1 x_2} Q b) \\ &\times \exp[-S(x_1, x_2, b, Q)] . \end{aligned} \quad (17)$$

where the complete Sudakov exponent is given by

$$S(x_1, x_2, b, Q) = \sum_{i=1}^2 [s(x_i, b, Q) + s(1 - x_i, b, Q)] + 4 \int_{1/b}^t \frac{d\bar{\mu}}{\bar{\mu}} \gamma_q(\alpha_s(\bar{\mu})) . \quad (18)$$

K_0 is the modified Bessel function of order zero, which comes from the Fourier transform to b space of the gluon propagator. Note that we insert the two-loop α_s in Eq. (10) into the above integral of γ_q , while it is the one-loop α_s that was employed in [3]. The exponential e^{-S} decreases rapidly at large b , leading to Sudakov suppression mentioned before. The essential advantage of Eq. (17), compared to Eq. (1), is then the extra b dependence in the hard scattering. If b is small, radiative corrections with the argument of α_s set to t will be small, regardless of the values of x . Of course, when b is large and $x_1 x_2 Q^2$ is small, radiative corrections are still large. However, we shall show that e^{-S} suppresses the contributions from this region, and the perturbation theory becomes relatively self-consistent.

In Eq. (17) the evolution of ϕ with $1/b$ is written as [21]

$$\phi(x, 1/b) = \frac{3f_\pi}{\sqrt{2N_c}} x(1-x) \left[\sum_{n=0}^{\infty} a_n C_n^{3/2}(2x-1) \left(\frac{\alpha_s(1/b)}{\alpha_s(\mu_0)} \right)^{\gamma_n} \right] , \quad (19)$$

where the coefficients a_n and the exponents γ_n are

$$a_n = \frac{5(2n+3)}{(n+2)(n+1)} I_n , \quad (20)$$

$$\gamma_n = \frac{4}{33-2n_f} \left[1 + 4 \sum_{k=2}^{n+1} \frac{1}{k} - \frac{2}{(n+1)(n+2)} \right] , \quad (21)$$

with $I_0 = 2/15$, $I_1 = 0$, $I_2 = 8/35$ and $I_n = 0$ or $n \geq 3$. The relevant Gegenbauer polynomials are $C_0^{3/2}(x) = 1$ and $C_2^{3/2}(x) = (15/2)(x^2 - 1/5)$. Then Eq. (19) reduces to

$$\phi(x, 1/b) = \frac{3}{\sqrt{2N_c}} f_\pi x(1-x) \left[1 + [5(1-2x)^2 - 1] \left(\frac{\alpha_s(\tilde{\mu})}{\alpha_s(\mu_0)} \right)^{\frac{50}{81}} \right] , \quad (22)$$

where we have changed the argument of α_s from $1/b$ to

$$\tilde{\mu} = \max(1/b, m) , \quad (23)$$

with m a mass scale greater than the initial value $\mu_0 = 0.5$ GeV. This modification, implying that $\phi(x, 1/b)$ is “frozen” at $\phi(x, m)$ as $1/b < m$, guarantees Eq. (22) to evolve toward the correct direction. Without this freezing scale, $\phi(x, 1/b)$ is ill-defined for $1/b \rightarrow \Lambda$, and large infrared contributions to the pion form factor will be observed. m will be treated as a parameter of order μ_0 , and determined by the data fitting. Obviously, Eq. (22) approaches the asymptotic wave function ϕ^{AS} in Eq. (3) as $1/b \rightarrow \infty$, and the preasymptotic wave function ϕ^{CZ} in Eq. (4) as $1/b \rightarrow \mu_0$. We then expect that the pion form factor derived from Eq. (22) is located at between those from ϕ^{AS} and ϕ^{CZ} .

We are also interested in the full expression of H in Eq. (15), except for the approximate one in Eq. (16). Without neglecting the virtual quark transverse momenta, the parallel formula to Eq. (17) involves a double- b integral:

$$\begin{aligned}
F_\pi(Q^2) &= 16\pi\mathcal{C}_F \int_0^1 dx_1 dx_2 \int_0^\infty b_1 db_1 b_2 db_2 \phi(x_1, 1/b_1) \phi(x_2, 1/b_2) \\
&\times \alpha_s(t) K_0(\sqrt{x_1 x_2} Q b_1) \exp[-S(x_1, x_2, b_1, b_2, Q)] \\
&\times [\theta(b_1 - b_2) K_0(\sqrt{x_1} Q b_1) I_0(\sqrt{x_1} Q b_2) + \\
&\quad \theta(b_2 - b_1) K_0(\sqrt{x_1} Q b_2) I_0(\sqrt{x_1} Q b_1)] , \tag{24}
\end{aligned}$$

with I_0 the modified Bessel function of order zero, and the complete Sudakov exponent

$$S = \sum_{i=1}^2 s(x_i, b_i, Q) + s(1 - x_i, b_i, Q) + 2 \int_{1/b_i}^t \frac{d\bar{\mu}}{\bar{\mu}} \gamma_q(\alpha_s(\bar{\mu})) , \tag{25}$$

and the hard scale

$$t = \max[\sqrt{x_1 x_2} Q, 1/b_1, 1/b_2] . \tag{26}$$

A freezing scale is not necessary for Eq. (24), since the extra degrees of freedom b_2 moderate the infrared enhancement from the low end of the evolution of the wave functions. This will be explicitly demonstrated in next Section.

III. Numerical Results

In the numerical analysis the exponential e^{-S} will be treated as follows. In the small b region higher-order corrections should be absorbed into the hard amplitude, instead of into the wave functions, giving their evolution in b . Hence, we set any Sudakov factor $e^{-s(\xi, b, Q)}$ to unity for $\xi Q/\sqrt{2} < 1/b$. In this vein, we also set e^{-S} to unity, whenever it exceeds unity in the small b region. As b increases, e^{-S} decreases, reaching zero at $b = 1/\Lambda$. Suppression in the large b region is weaker for smaller Q , where Sudakov effects are mild. The above typical behavior of e^{-S} has been shown in [3]. It is expected that the suppression improves the self-consistency of the perturbative calculation of the pion form factor.

We evaluate $Q^2 F_\pi(Q^2)$ employing the pion wave function ϕ with evolution in Eq. (22). We find, at $Q^2 = 9 \text{ GeV}^2$, $Q^2 F_\pi(Q^2) = 0.33 \text{ GeV}^2$ for the freezing scale $m = \mu_0$ and $Q^2 F_\pi(Q^2) = 0.28 \text{ GeV}^2$ for $m = 2\mu_0$, the latter being only 15% smaller than the former. It implies that our formalism is insensitive to the choice of m . If a freezing scale was not introduced, the predictions are almost enlarged by a factor 6, which is attributed to the infrared enhancement from the region with $1/b < \mu_0$, *i.e.*, to the misuse of Eq. (22). We adopt the natural choice $m = \mu_0$ in the analysis below.

As to the self-consistency of the perturbative calculation, we propose to analyze Eq. (17), not by cutting off the x integrals near their endpoints [22], but by testing the sensitivity of the integral to the large b region. We expect that, because Eq. (1) is the true asymptotic behavior, as Q increases, the b integral will become more and more dominated by the small b contributions, for which the effective coupling is small. To see how the contributions to Eq. (17) are distributed in b space under Sudakov suppression, the integration is done with a variable cut off in b , b_c . Typical numerical results are displayed in Fig. 3 for the use of ϕ in Eq. (22) with $\Lambda = 0.15 \text{ GeV}$. The curves, showing the dependence of $Q^2 F_\pi(Q^2)$ on b_c , rise from zero at $b_c = 0$, and reach their full height at $b_c = 1/\Lambda$, beyond which we consider any remaining contributions as being truly nonperturbative. As anticipated, we observe a faster rise as Q increases.

To be quantitative, we may consider the cutoff up to which half of the whole contribution has been accumulated. For $Q^2 = 4$ and 9 GeV^2 , 50% of $Q^2 F_\pi(Q^2)$ comes from the regions with $b \leq 4.0 \text{ GeV}^{-1}$ [$\alpha_s(1/b)/\pi \leq 0.44$] and with $b \leq 3.2 \text{ GeV}^{-1}$ [$\alpha_s(1/b)/\pi \leq 0.37$], respectively. A liberal standard to judge the relevance of the perturbative method would be that 50% of the

results come from the region where $\alpha_s(1/b)/\pi$ is no larger than, say, 0.5. From this point of view, PQCD begins to be self-consistent at $Q^2 \sim 4 \text{ GeV}^2$. This conclusion is consistent with that drawn in [3], where the CZ wave function ϕ^{CZ} and the asymptotic wave function ϕ^{AS} were employed.

Results of $Q^2 F_\pi(Q^2)$ for the use of ϕ^{AS} , ϕ^{CZ} and ϕ in Eq. (22), as well as the experimental data [6, 7, 8, 9], are shown in Fig. 4. As expected, the curve corresponding to Eq. (22) is located at between the other two. It is known that the pion form factor behaves like $1/Q^2$ asymptotically. The data points of $Q^2 F_\pi(Q^2)$, arising from low Q^2 and reaching a plateau at $Q^2 \sim 2 \text{ GeV}^2$, reflect this asymptotic behavior. Hence, the point at $Q^2 \sim 10 \text{ GeV}^2$, jumping to a value about twice of the plateau, seems not plausible, and is not considered here. In this sense, our predictions are in a good agreement with the data for $Q^2 > 4 \text{ GeV}^2$.

For $Q^2 < 4 \text{ GeV}^2$, all the curves rise rapidly and deviate from the data, indicating the dominance of the nonperturbative contrubutions. Note that the predictions from ϕ^{AS} and ϕ^{CZ} are larger than those obtained in [3]. This is due to the insertion of the two-loop α_s into the integral of γ_q in Eq. (18). To examine the stability of our predictions to the variation of the scales Λ and μ_0 , we compute $Q^2 F_\pi(Q^2)$ for $\Lambda = 0.1$ and 0.2 GeV ($\mu_0 = 0.5 \text{ GeV}$), and for $\mu_0 = 0.4$ and 0.6 GeV ($\Lambda = 0.15 \text{ GeV}$) at $Q^2 = 9 \text{ GeV}^2$. Their values are listed in Table I. Apparently, the difference from those corresponding to $\Lambda = 0.15 \text{ GeV}$ and $\mu_0 = 0.5 \text{ GeV}$ is small.

At last, we evaluate $Q^2 F_\pi(Q^2)$ using the double- b factorization formula in Eq. (24), and the results are presented in Fig. 5. Those derived from the single- b formula in Eq. (17) and the data are also displayed for comparision. We find that the two curves are close to each other especially in the region with large $Q^2 \sim 5\text{-}10 \text{ GeV}^2$. The consistency of the predictions from Eq. (24) with the data is nontrivial, since it does not involve an adjustable freezing scale m . The above analyses indicate that our scheme of truncating the infrared enhancement from the evolution of the wave function indeed makes sense. In the future study of the proton form factor, we shall take the same approach in order to simply the analysis, that is, neglect the transverse momenta carried by the virtual quarks, and introduce a freezing scale for the proton wave function.

IV. Conclusion

In this paper we have performed a very thorough investigation of the pion form factor based on the modified PQCD formalism. Both the two approaches with the single- b formula plus the introduction of a freezing scale and with the double- b formula lead to results that match the experimental data. We have also confirmed that our predictions are dominated by perturbative contributions for $Q^2 > 4 \text{ GeV}^2$, and insensitive to the variation of the parameters Λ and μ_0 . All these conclusions imply that our formalism is reliable for the study of exclusive QCD reactions at intermediate energy scales. We shall extend it to more complicated processes, such as the proton form factor and Compton scattering, which will be published elsewhere.

References

- [1] N. Isgur and C.H. Llewellyn Smith, Nucl. Phys. B317 (1989) 526.
- [2] T. Huang and Q.X. Shen, Z. Phys. C50 (1991) 139.
- [3] H-n. Li and G. Sterman, Nucl. Phys. B381 (1992) 129.
- [4] G.P. Lepage and S.J. Brodsky, Phys. Rev. Lett. 43 (1979) 545; Phys. Rev. D22 (1980) 2157.
- [5] G.P. Lepage and S.J. Brodsky, Phys. Lett. B87 (1979) 359.
- [6] C.N. Brown *et al.*, Phys. Rev. D8 (1973) 92.
- [7] C.J. Bebek *et al.*, Phys. Rev. D9 (1974) 1229.
- [8] C.J. Bebek *et al.*, Phys. Rev. D13 (1976) 25.
- [9] C.J. Bebek *et al.*, Phys. Rev. D17 (1978) 1693.
- [10] V.L. Chernyak and A.R. Zhitnitsky, Nucl. Phys. B201 (1982) 492; Phys. Rep. 112 (1984) 173; Nucl. Phys. B246 (1984) 52.
- [11] N. Isgur and C.H. Llewellyn Smith, Phys. Rev. Lett. 52 (1984) 1080.
- [12] A.V. Radyushkin, Acta Phys. Polonica, B15 (1984) 403; A.P. Bakulev and A.P. Radyushkin, Phys. Lett. B271 (1991) 223.
- [13] J.C. Collins and D.E. Soper, Nucl. Phys. B193, 381 (1981).
- [14] J. Botts and G. Sterman, Nucl. Phys. B325 (1989) 62
- [15] J.C. Collins, in *Perturbative Quantum Chromodynamics*, A.H. Mueller ed. (World Scientific, Singapore, 1989).
- [16] G.R. Farrar and D.R. Jackson, Phys. Rev. Lett. 43 (1979) 246
- [17] A.V. Efremov and A.V. Radyushkin, Phys. Lett. B94 (1980) 245.
- [18] V.L. Chernyak and A.R. Zhitnitsky, Yad. Fiz. 31 (1980) 1053 [Sov. J. Nucl. Phys. 31 (1980) 544; Nucl. Phys. B216 (1983) 373.

- [19] C.E. Carlson and F. Gross, Phys. Rev. D36 (1987) 2060.
- [20] R. Jakob and P. Kroll, Phys. Lett. B315 (1993) 463. J. Bolz, R. Jakob, P. Kroll, M. Bergmann and N.G. Stefanis, Z. Phys. C66 (1995) 267.
- [21] C.R. Ji and F. Amiri, Phys. Rev. D42 (1990) 3764.
- [22] A.V. Radyushkin, Nucl. Phys. A532 (1991) 141.

Table I. The values of $Q^2 F_\pi(Q^2)$ for different Λ and μ_0 at $Q^2 = 9 \text{ GeV}^2$.

$\Lambda(\text{GeV})$	$Q^2 F_\pi(Q^2)(\text{GeV}^2)$	$\mu_0(\text{GeV})$	$Q^2 F_\pi(Q^2)(\text{GeV}^2)$
0.10	0.2974	0.4	0.3114
0.15	0.3309	0.5	0.3309
0.20	0.3659	0.6	0.3333

Figure Captions

Fig. 1. (a) Factorization of the pion form factor. The symbol $\times$ represents the photon vertex. (b) Basic scattering diagrams.

Fig. 2. Radiative corrections to the basic scattering diagrams.

Fig. 3. Dependence of $Q^2 F_\pi(Q^2)$ on b_c for the use of ϕ in Eq. (22) at $Q^2 = 4$ GeV^2 (dashed line) and at $Q^2 = 9$ GeV^2 (solid line).

Fig. 4. Dependence of $Q^2 F_\pi(Q^2)$ on Q^2 for the use of ϕ^{AS} (dotted line), of ϕ^{CZ} (dashed line), and of ϕ in Eq. (22) (solid line).

Fig. 5. Dependence of $Q^2 F_\pi(Q^2)$ on Q^2 for the use of ϕ in Eq. (22) derived from Eq. (17) (dashed line) and from Eq. (24) (solid line).

This figure "fig1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9712541v1>

This figure "fig2.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9712541v1>

This figure "fig3.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9712541v1>

This figure "fig4.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9712541v1>

This figure "fig5.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9712541v1>